

# A Pure Mathematical Proof of the 4-Colour Theorem

**Jun Wang\***

*Master Of Condensed Matter Physics of Sun Yat-Sen University, China*

**\*Corresponding author:** Jun Wang, Master of Condensed Matter Physics of Sun Yat-Sen University, China

## ARTICLE INFO

**Received:** 📅 September 06, 2023

**Published:** 📅 September 19, 2023

**Citation:** Jun Wang. A Pure Mathematical Proof of the 4-Colour Theorem. Biomed J Sci & Tech Res 52(5)-2023. BJSTR. MS.ID.008323.

## ABSTRACT










This work "A Pure Mathematical Proof of the 4-Colour Theorem" is related to the previous proof assisted by computer. "Triangulations of Euler Convex Polygon" provides a fresh beginning point for the proof. The central concept is to discover an extended invariance property of Standard Graph's boundary, which is described as "3-Colour All Phase States (3CP)" in this work and it is demonstrated that the Standard Graph's boundary and sub-bound are 3CP and 4-colorable (4-3CP) via the Expanded Operation  $e(+, \pi)$  and  $e(-, \pi)$ . It's exciting that this regularity was discovered for the first time and the 4-3CP invariance can naturally derive the 4-Colour Theorem. The majority of the definitions, theorems and proof strategies are shown in this work.

**Keywords:** 4-Colour Theorem; Standard Graph; Bound; Sub-bound; 3CP;  $e(+, \pi)$ ;  $e(-, \pi)$ ; Triangulations

## Introduction

The 4-Colour Theorem states that every map can be coloured using only four colours and no two adjacent regions have the same colour. The initial problem was first posed in the mid-19<sup>th</sup> century by Francis Guthrie. Guthrie noticed that four colours were sufficient to colour the map, and he wondered if this was true for every map. Many mathematicians attempted to solve the problem, but rigorous mathematical proof remained elusive. It became one of the most famous mathematical problems of the 20<sup>th</sup> century and Kenneth Appel and Wolfgang Haken finally solved computer-aided proof in 1976. This proof was considered controversial due to the extent of the computer assistance required. The 4-Colour Theorem has important applications in real-world situations, such as in scheduling and timetabling problems. It also demonstrates the power and elegance of mathematical reasoning, as well as the importance of collaboration and innovation in solving complex problems. We describe a specific class of graphs (Appendix Figure 1) called Standard Graphs (SG), which are constructed by the Expanded Operations  $e(+, p_i)$  and  $e(-, p_i)$ , which

are used to replace planar graphs since they may have a significant number of unsaturated links and lack uniformity. The Proof begins with "Triangulations of Euler Convex Polygons": convex polygons can be sliced into multiple triangles (Euler obtains the famous Catalan Number [1] by counting the number of triangulations of convex polygons [2]). In this article, we defined all the possible triangulations sets of convex polygons (denoted by "bound") as "All Phase States (AP)". If the boundary of any Standard Graph (SG) is All Phase States (AP) and 4-colorable, the 4-Colour Theorem will be established [3]. Unfortunately, the All Phase States (AP) property of the boundary cannot keep Invariance during the Expanded Operation  $e(+, p_i)$  (denoted by "IEO"), which implies that the All Phase States (AP) property is incomplete. Thus the boundary of the Standard Graph must have a cryptic property that can satisfy both the All Phase States (AP) and IEO. Finally, I find that the Standard Graph's boundary satisfy 3CP and 4-colorable (denoted by "4-3CP") and it is proven 4-3CP is IEO. This article's primary goal is to attest the Standard Graph's boundary  $J(m)$  and all its sub-bounds set  $\{J(m')\}$  are 4-3CP.

Definition	Samples	Definition Description
Point (P)	$\cdot p$	A country or area in a graph, $P(G)$ is called point set of graph(G).
Link (l)	$p_1 \leftrightarrow p_2$ 	The relationship between two adjacent points $\{p_1, p_2\}$ , $L(G)$ is called link set of graph(G).
Element (e)		For the point set $P = \{a, d, e\}$ , a, d, e link with each other form a $K^3$ , we call it is an <b>element e(ade)</b> , $E(G)$ is elements set of graph(G).
Bound $J(m)$		Set a graph $G(P_m, L_m)$ , $P_m = \{p_1, p_2, \dots, p_m\}$ , $L_m = \{p_1p_2, p_2p_3, \dots, p_{m-1}p_m, p_m p_1\}$ , $m \geq 3$ , where the points $p_1, p_2, \dots, p_m$ are all distinct and form a cycle $C^m$ , then the cycle $C^m$ is called a <b>bound J(m) or J(P_m)</b> . $ J(m)  = m$ is number of points of $J(m)$ , $\ J(m)\  = m$ is number of links of $J(m)$ .
Sub-bound $J(m')$ / Com-bound $\overline{J(m')}$		Set a bound $J(P_m)$ , we get points set $P_{m'} \subseteq P_m$ form a new bound $J(P_{m'})$ or $J(m')$ , which is called <b>sub-bound</b> of $J(P_m)$ . We call $J(m')$ , $J_1, J_2, \dots$ (without cut links and loops) is <b>sub-bound set <math>\{J(m')\}</math></b> of $J(P_m)$ . The bound set $\{J_1, J_2, \dots\}$ is called <b>com-bound <math>\overline{J(m')} = J(m) - J(m') = J_1 + J_2 + \dots</math></b> of $J(m')$ in $J(m)$ .
$e(+, p_i)$		The operation of adding a element $e(ade)$ to the bound $J(P_m)$ to become a new bound $J(m+a)$ or $J(P_m+a)$ ( $d, e \in P_m, d-e$ ). $ J(P_m+a)  -  J(P_m)  = 1$ . We denote this bound expanded operation as <b><math>e(+, p_i)</math></b> , $p_i = d \in P_m$
$e(-, p_i)$		The operation of removing a point a from the bound $J(P_m)$ to become a new bound $J(m-a)$ or $J(P_m-a)$ ( $a, d, e \in P_m, d-a, a-e$ ). $ J(P_m-a)  -  J(P_m)  = -1$ . We denote this bound expanded operation as <b><math>e(-, p_i)</math></b> , $p_i = d \in P_m$
Standard Graph SG/ Subgraph SG'		The graphs constructed by $e(+, p_i)$ and $e(-, p_i)$ is <b>standard graph (SG)</b> , <b><math>SG(n) = \sum_{i=1}^n e(\pm, p_i)</math></b> , $n$ is the number of $e$ . we call $Y$ is the colouring solution set family of boundary $J(m)$ of $SG(n)$ , $ Y $ is chromatic number of $J(m)$ , the 3-colouring solution set of $Y$ denote by $3Y$ , $ 3Y  = 3$ . $m =  e(+, p_i)  -  e(-, p_i)  + 2$ ; The standard graph SG removed some links (red links) is <b>subgraph <math>SG(n')</math></b> .
All Phase States (AP)		All link states without cut link of the bound $J(m)$ are called <b>all phase states (AP)</b> . The all phase states number of $J(m+2)$ is $\frac{1}{m+1}C_{2m}^m = C(m)$ , ( $C(m)$ is Catalan Number)
3-Colour All Phase States (3CP)		Let $J(P_m)$ be a bound, $Y$ is the colouring solution set family of $J(P_m)$ , $\exists  Y  \leq 4$ , $Y$ must have $3Y$ and $Y$ can make the bound $J(P_m)$ be AP, we call $J(P_m)$ is <b>3-Colour All Phase State (3CP)</b> .

Appendix Figure 1: Definition

### Bound

Set a cycle  $C(P_m, L_m)$ ,  $P_m = \{p_1, p_2, \dots, p_m\}$ ,  $L_m = \{p_1p_2, p_2p_3, \dots, p_{m-1}p_m, p_m p_1\}$ ,  $m \geq 3$ , cycle  $C$  divides the plane into two connected domain: inside and outside, then we call the cycle  $C$  be a bound (Figure 1), denoted by  $J(m)$  or  $J(P_m)$ .  $|J(m)| = m$  is the number of points of  $J(m)$ ,  $||J(m)|| = m$  is the number of links of  $J(m)$ . There are two types of link: real link “—” ( $p_1p_2, p_2p_3, \dots$ ) and virtual link “...” ( $p_1 \dots p_4, p_3 \dots p_5, \dots$ )

although the two points aren't linked by a real link, the virtual link is exist between them cause they are colored differently. Set a bound  $J(m)$ , we get points set  $P_{m'} \subseteq P_m$  to form some new bounds  $J(m'), J_1, J_2, \dots$ , (without cut links and loops), which are called sub-bounds of  $J(m)$ . The  $J(m'), J_1, J_2, \dots$  is sub-bounds set  $\{J(m')\}$  of  $J(m)$ . The sub-bounds set  $\{J_1, J_2, \dots\}$  is called com-bound  $\bar{J}(m') = J(m) - J(m') = J_1 + J_2 + \dots$  of  $J(m')$  in  $J(m)$ .




Figure 1: Bound.

### All Phase States

We defined a triangulation of convex polygons as a link state and provided samples of  $J(3)$ - $J(7)$  in Figure 2. If the bound  $J(m)$  contains all link states, we call  $J(m)$  all phase states (AP). The all link states number of  $J(m+2)$  is  $C(m) = \frac{1}{m+1} C_{2m}^m$ , ( $C(m)$  is Catalan Number). Set

$\{J(m')\}$  be a sub-bound set of  $J(m)$ , it is simple to demonstrate that all of the sub-bounds set  $\{J(m')\}$  are AP if and only if  $J(m)$  is AP. The sub-bounds set  $\{J(m')\}$  is AP, which means that the sub-bound  $J(m') \in \{J(m')\}$  is AP and independent.




Figure 2: Link states of  $J(3)$ - $J(7)$ .

### The 3-Colour All Phase States (3CP)

Let  $J(P_m)$  be the boundary of Standard Graph SG (Figure 3),  $Y$  is the colouring solution set family of  $J(P_m)$ , if  $Y$  can make the bound  $J(P_m)$  be AP and  $\exists 3Y \in Y$ , we call  $J(P_m)$  3-Colour All Phase State (3CP). An

example of 3CP is provided bellow (Figure 4). If we have enough time to test all of the Standard Graph conditions, we will discover that the boundary of every Standard Graph is 3CP and 4-colorable (Appendix Text). It's exciting that this regularity was discovered for the first time and the 4-3CP invariance can naturally derive the 4-Colour Theorem.




Figure 3: An example of the 3-Colour All Phase States (3CP).



Figure 4:  $J(P_m)$  of SG.

### The 4-3CP Conjecture

The conjecture is described as follows:

4-3CP Conjecture:  $\forall$  standard graph SG,  $J(m)$  is the boundary of SG, let  $Y$  is the colouring solution set family of  $J(m)$ ,  $\{J'(m)\}$  is sub-bounds set of  $J(m)$ ,  $Y$  can make:


- (1)  $|Y| \leq 4$ ,
- (2)  $J(m)$  be 3CP,
- (3)  $\{J'(m)\}$  be 3CP,
- (4)  $\exists a 3-colour solution set 3Y(J'(m))$  and  $\{\bar{J}'(m)\}$  be 3CP,
- (5) Let  $x, p_1, p_2 \in J'(m)$ , link  $x-p_1, x-p_2$  form sub-bounds  $J_1(x-p_1), J_2(x-p_2)$ , and  $\overline{J_1+J_2}$ , if  $x$  scans on  $p_1 \rightarrow p_2, \exists 3Y(J_1+J_2)$  and  $\overline{J_1+J_2}$  is 3CP.

### Proof

By Exhaustive Method, It's Easy to See

- (1) Element  $e = SG(1), J(3)$  conform to 4-3CP Conjecture.
- (2)  $SG(2), J(4)$  conform to 4-3CP Conjecture.
- (3)  $SG(3), J(3)$  conform to 4-3CP Conjecture.

Set a Standard Graph SG, the boundary  $J(P_m)$  of SG, the colouring solution set family  $Y$  of  $J(P_m), |Y| \leq 4, J(P_m)$  and its sub-bounds  $J(P_m')$  conform to the 4-3CP Conjecture. Then add an element  $e(ade)$  (points  $a, d, e$  form a triangle) on the  $J(P_m)$  to form a new boundary denoted by  $J(a + P_m) = J(P_m) + e(ade)$  ( $d, e \in J(P_m), d-e, a$  is new). According to the Points Scanning Method (Appendix Figure 3), it's easy to prove  $J(a + P_m)$  is 3CP and 4-colorable, so 4-3CP Conjecture (2) is proven. Next, we shall prove the sub-bounds set  $\{J(a + P_m')\}$  of  $J(a + P_m)$  is 3CP and 4-colorable: When  $|Y| = 3, J(P_m)$  is AP equals to  $J(P_m)$  is 3CP. Since  $J(P_m)$  is AP, so the sub-bounds set  $\{J(P_m')\}$  is AP, and  $Y(J(P_m')) \in Y, |Y(J(P_m'))| = |Y| = 3$ , so  $\{J(P_m')\}$  is 3CP. Then we shall prove when  $|Y| = 4$ , the sub-bounds set  $\{J(a + P_m')\}$  of  $J(a + P_m)$  is 3CP.



Appendix Figure 3:  $J(m)$  Links Scanning Method.

**When the New Point A  $\notin$  3-Colour Sub-Bound Of  $J(A+P_m)$**

Take any  $n$  points  $P_n = \{p_1, p_2, p_3, \dots, p_n\} \subseteq P_m$ , link  $p_1-p_2, p_2-p_3, \dots, p_{n-1}-p_n$ , form  $(n-1)$  sub-bounds  $J(p_1-p_2), J(p_2-p_3), \dots, J(p_{n-1}-p_n)$ , and take any point  $x, x \in P_n, x \neq p_1, p_n$ , link  $x$  with  $p_1, p_n$  form two sub-bounds  $J(x \rightarrow p_1)$  and  $J(p_1 \rightarrow x)$ ; link  $x$  with  $d, e$  form  $J(x \rightarrow p_n d), J(x \rightarrow ep_1)$  and  $e(xde)$  (Figure 5). According to  $J(P_m)$  and the sub bounds of  $J(P_m)$  are 4-3CP:  $J(p_1-p_2), J(p_2-p_3), \dots, J(p_{n-1}-p_n)$  and  $J(x \rightarrow p_n), J(p_1 \rightarrow x)$  and  $J(x \rightarrow p_n d), J(x \rightarrow ep_1), e(xde)$  are 3CP. According to 4-3CP Conjecture, when  $x$  scans on  $P_n (x \neq p_1, p_n)$ , the colouring solution set family  $Y(P_n) (Y(P_n) \in Y)$  must have a 3-colouring solution set  $3Y(P_n)$  make  $J(p_1-p_2), J(p_2-p_3), \dots, J(p_{n-1}-p_n)$  and  $J(x \rightarrow p_n d), J(x \rightarrow ep_1), e(xde)$  are 3CP. If  $y(a) = y(x), 3Y(P_n)$  will make  $|Y(J(a + P_n))| = 3$  and make  $J(a \rightarrow p_n d), J(a \rightarrow ep_1)$  and  $J(p_1-p_2), J(p_2-p_3), \dots, J(p_{n-1}-p_n)$  are 3CP, which means  $\{J(a + P_n)\}$  are 3CP. So,  $\exists a$  3-colouring solution set  $3Y(J(a + P_n))$  make  $\{J(a + P_n)\}$  be 3CP. 4-3CP Conjecture (4) is proven.




Figure 5.

**When The New Point A  $\notin$  3-Colour Sub-Bound Of  $J(A + P_m)$**

Since point  $a \notin$  3-colour sub-bound of  $J(a + P_m)$ , we shall to prove all sub-bound  $J(a + P_m')$  with point  $a$  are AP. Take any  $n + 2$  points  $P_n = \{d, e, p_1, p_2, p_3, \dots, p_n\} \subseteq P_m$ , link  $e-p_1, p_1-p_2, p_2-p_3, \dots, p_{n-1}-p_n, p_n-d$ , form  $(n + 2)$  sub-bounds  $J(p_1-p_2), J(e-p_1), J(p_2-p_3), \dots, J(p_{n-1}-p_n), J(p_n-d)$  and  $J(a + P_n)$ . (Figure 6) According to the boundary  $J(P_m)$  and the sub bounds of  $J(P_m)$  are 3CP:  $J(P_n), J(e-p_1), J(p_1-p_2), J(p_2-p_3), \dots, J(p_{n-1}-p_n), J(p_n-d)$  are 3CP,  $\exists$  3-colouring solution set  $3Y(J(P_n)) \in Y$ , make  $\{J(P_n)\}$  be 3CP. According to the Points Scanning Method: If  $y(a) \notin 3Y(J(P_n))$  and  $y(a) \in Y, J(a + P_n)$  is AP, so all sub-bound  $J(a + P_m')$  with point  $a$  are AP. Sum up 5.1.1, 5.1.2, we can see 4-3CP Conjecture (3) is proven.

**Proof of 4-3CP Conjecture (5)**

Set  $P_m' \subseteq P_m, P_m'$  divide  $J(m)$  into  $m'$  parts, let  $x$  is a point between  $p_1 \rightarrow p_2 (x \neq p_1, p_2)$  on  $J(m)$ , we call  $p_1, p_2$  are fixed points,  $x$  is scanning point. Link  $x-p_1, x-p_2, \dots, x-p_{m'}$ , form a sub-bounds set  $\{J_i(x) | x \in J_i(x), i=1, 2, \dots, m'\}$  of  $J(m)$ . (assume  $m' = 5$ , see Figure 7). According to 4-3CP Conjecture (5), when  $x$  scans on  $p_1 \rightarrow p_2, \exists 3Y(J_1 + J_2)$ , and

$J_3, \dots, J_m', J_{m'+1}$  are 3CP. Then add a new point  $a$  on  $J(m)$ , when point  $a$  is on  $J_1, J_2, \exists y(a) \in Y(J_1(x) + J_2(x))$ , keep  $|Y(J_1(x) + J_2(x))| = 3$ , and  $J_3, \dots, J_m', J_{m'+1}$  are 3CP; when point  $a$  is on  $J_3, \dots, J_m', J_{m'+1}, \exists y(a) \in Y$ , keep  $J_3, \dots, J_m', J_{m'+1}$  are 3CP; when point  $a$  is a fixed point as  $p_1, p_2$ , we also could prove that, when  $x$  scans on  $p_1 \rightarrow p_2, \exists 3Y(J_1 + J_2)$ , and  $J_3, \dots, J_m', J_{m'+1}$  are 3CP (we will prove in detail in next article), so  $J(m + a)$  can keep when  $x$  scans on  $p_1 \rightarrow p_2, \exists 3Y(J_1 + J_2)$ , and  $J_3, \dots, J_m', J_{m'+1}$  are 3CP. So 4-3CP Conjecture (5) is proven (Figure 8).




Figure 6: a E 3-colour sub-bound of  $J(a + P_m)$ ,  $a = x$ .




Figure 7: a  $\notin$  3-colour sub-bound of  $J(a + P_m)$ .




Figure 8: Sub-bounds set  $\{J_i(x) \mid x \in J_i(x), J_i(x) \in J_1, \dots, J_6\}$  of  $J(m)$ .

So  $Y(a + m)$  can make:

- (1) So  $|Y(a + m)| = |Y| \leq 4$ ,
- (2) The bound  $J(a + m)$  be 3CP,
- (3)  $\forall$  sub-bounds set  $\{J'(a + m)\}$  be 3CP,
- (4)  $\exists$  a 3-colour solution set  $3Y(J'(a + m))$ , and  $\{\bar{u}(+)\}$  be 3CP.
- (5) Let  $x, p_1, p_2 \in J'(a+m)$ , link  $x-p_1, x-p_2$  form sub-bounds  $J'_1(x-p_1), J'_2(x-p_2)$ ,  $J'_1 + J'_2$ , if  $x$  scans on  $p_1 \rightarrow p_2$ ,  $\exists 3Y(J'_1 + J'_2)$  and  $J'_1 + J'_2$  is 3CP. Above all, we have proven 4-3CP Invariance during the Expanded Operation  $e(+, p_i)$ , It's easy to see 4-3CP Invariance during the Expanded Operation  $e(-, p_i)$  is also true. So 4-3CP Conjecture is proven!

### Conclusion

The major research objects are the boundary's invariance property of Standard Graph in this work. By creating the Standard Graph via Expanded Operations  $e(+, p_i)$  and  $e(-, p_i)$ , the complexity of the planar graph is reduced. By examining the Invariance during Expanded Operations (IEO) of Standard Graph, huge calculation for coloring planar graph are avoided. Based on the above optimization method, we have demonstrated a rigorous proof of the 4-Colour Theorem, which can be also used to optimize complex systems..

### References

1. Jun Wang, Hisham MN, Al Selwi MK, Al-Shwafi AN (2018) Assessment of Heavy Metals Concentration in Mangroves Leaves of the Red Sea Coast of Yemen. *J Ecol & Nat Resour* 2(1): 120.
2. Shruthi, Padmanabha A, Lakshmi pathi MT (2022) Diatom diversity in relation to physico-chemical parameters of coastal waters of Mangalore and Padubidri, Southwest coast of India. *International Journal of Ecology and Environmental Sciences* 4(4): 28-37.

3. Sagar M, Waghmare, Pramod T, Hanamgond, Debashish Mitra, et al. (2020) Application of Remote sensing and GIS techniques to study sediment movement along Harwada Beach, Uttar Kannada, West coast of India. *Journal of Coastal Research* 36(6): 1121-1129.
4. Karande AA (1991) Use of epifaunal communities in pollution monitoring. *J Environ Biol* 12: 191- 200.
5. Poonam Bhadja, Rahul Kundu (2012) Status of seawater quality at few industrially important coasts of Gujarat (India) off Arabian Sea. *Indian Journal of Geo-Marine Sciences* 41(1): 90-97.
6. GJ Chakrapani, V Subramanian (1993) "Rate of Erosion and Sedimentation in the Mahanadi River Basin, India". *Journal of Hydrology* 149(1-4): 39-48.
7. Soren Laurentius Nielsen, Gary T Banta, Morten Foldager Pedersen (2004) "Estuarine Nutrient Cycling: The Influence of Primary Producers", Kluwer Academic Publishers, the Netherlands.
8. Sridhar R, Thangaradjou T, Kannan L (2008) Comparative investigation on physico-chemical properties of the coral reef and seagrass ecosystems of the Palk Bay. *Indian Journal of Mar. Sciences* 37(2): 207-213.
9. Asha PS, Diwakar (2007) Hydrobiology of the inshore waters off Tuticorin in the Gulf. *J. Mar. Biol. Ass. India* 49(1): 7-11.
10. Choudhury SB, RC Panigrahy (1991) Seasonal distribution and behavior of nutrients in the creek and coastal waters of Gopalpur, east coast of India. *Mahasagar-Bull. Natl. Inst. Oceanogr* 24: 81-88.
11. (2017) AHA. Standard methods for examination of water and wastewater. 23rd Edition, American Public Health Association, Washington, D.C. (USA), pp. 1200.
12. Jun Wang, Hisham MN, Al Selwi MK, Al Shwafi AN (2014). Hydrographical Studies on Mangroves Ecosystem of The Red Sea Coast of Yemen from Al-Salif to Bab-el-Mndab Strat. University of Aden *Journal of Natural and Applied* 18(2): 381-391.
13. Jha DK, Devi MP, Vidyalakshmi R, Brindha B, Vinithkumar NV, et al. (2015) Water quality assessment using water quality index and geographical information system methods in the coastal waters of Andaman Sea, India. *Mar Pollut Bull* 100(1): 555-561.
14. Abowei J (2010) Salinity, dissolved oxygen, pH and surface water temperature conditions in Nkoro River, Niger Delta, Nigeria. *Adv J Food Sci Technol* 2(1): 36-40.
15. Jun Wang (2017) Study of Some Heavy Metals Concentrations in Mangroves Environment - Red Sea Coast of Yemen. Unpublished M.Sc. Thesis, Fac Sci, Sana'a Univ, Yemen, pp. 161.
16. Ball MC (1988) Salinity Tolerance in the Mangroves *Aegiceras comiculatum* and *Avicennia marina*. I. Water use in relation to growth, carbon partitioning and salt balance. *Australian J Plant Physiol* 15(3): 447-464.
17. Suski C, Killen S, Keiffer J, Tufts B (2006) The influence of environmental temperature and oxygen concentration on the recovery of largemouth bass from exercise. Implications for live-release angling tournaments. *J Fish Biol* 68(1): 120-136.
18. Elahi N, Ahmed Q, Bat L, Yousuf F (2015) Physicochemical parameters and seasonal variation of coastal water from Balochistan coast, Pakistan. *J Coast Life Med* 3(3): 199-203.

ISSN: 2574-1241

DOI: 10.26717/BJSTR.2023.52.008323

Jun Wang. Biomed J Sci & Tech Res



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>



#### Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>