

Histological Profile of Primary Malignant Ovarian Tumors

Volahasina Francine Ranaivomanana, Holy Tiana Andrianjafitrimo*, Haingo Voahangy Rabetafika Ranaivoson, Tafitaherisoa Manajatiana Ravelomampitoninainarivony and Nantenaina Soa Randrianjafisamindrakotroka



Médecin Spécialiste et ancien interne en Anatomie et Cytologie Pathologiques, CHU-JRA Ampefiloha, Antananarivo, Madagascar

***Corresponding author:** Holy Tiana Andrianjafitrimo, Médecin Spécialiste et ancien interne en Anatomie et Cytologie Pathologiques, CHU-JRA Ampefiloha, Antananarivo, Madagascar

ARTICLE INFO

Received: 📅 July 11, 2022

Published: 📅 August 03, 2022

Citation: Volahasina Francine Ranaivomanana, Holy Tiana Andrianjafitrimo, Haingo Voahangy Rabetafika Ranaivoson, et al. Histological Profile of Primary Malignant Ovarian Tumors. Biomed J Sci & Tech Res 45(3)-2022. BJSTR. MS.ID.007212.

SUMMARY

Malignant germ cell tumors of the ovary are rare. They mainly affect young patients of childbearing age and can threaten the functional prognosis. We carried out a retrospective study of 11 cases of malignant germ cell tumors of the ovary, diagnosed at the Paraclinical Training and Research Unit in Anatomy and Pathological Cytology, over a period of 10 years, from January 2010 to December 2019. The age of the patients varied from 2 to 63 years with an average of 30, 45 years. In the majority of cases, radical treatment was performed with only one case of cystectomy and one case of tumor biopsy. Clinical signs were abdominopelvic mass in all cases. The most common histologic type was dysgerminoma (4 cases / 11). In our series, despite the young age of the majority of our patients, the treatment received was radical in 82% of cases, and therefore non-conservative.

Introduction

Primary malignant germ cell tumors of the ovary are rare. They represent 2 to 3% of ovarian tumors and mainly affect young patients, in the second and third decades of life [1]. The objective of our work was to determine the epidemiological and anatomopathological characteristics of these rare tumors.

Materials and Methods

We carried out a retrospective and descriptive study of the cases of primary malignant germ cell tumors of the ovary diagnosed at the paraclinical training and research unit in Pathological Anatomy and Cytology of the HU/JRA, over a period of 10 years, going from January 2010 to December 2019. We included all cases of malignant germ cell tumors of the ovary of any age. Our data collection

concerned the age, the affected side, the type of sample, the signs of clinical orientation, and the histological types.

Results

During the study period, we identified 68 patients in whom the diagnosis of malignant ovarian tumor was made. Of these cases, 11 were primary malignant ovarian germ cell tumors, accounting for 16.17%. The age of the patients varied from 2 to 63 years with an average of 30.45 years, but in 63.63% of cases, the patients were less than 25 years old. In the majority of cases (7 cases), an oophorectomy was performed. The right side was the most affected (6/11) and one case was bilateral. The guiding clinical information was in all cases an abdominopelvic mass isolated or associated with other signs (abdominal pain, weight loss) (Table 1).

Table 1: Epidemiological-clinical and anatomic-pathological characteristics of our patients.

N°	Age (Year)	Type of Direct Debit	Side	Macroscopy	Cut (Centimeter)	Diagnostic
1	2	Oophorectomy	Right	cerebral and cystic mass	19	Immature teratoma
2	21	Oophorectomy	Unspecified	Multiple cerebral fragments with cystic cavity	32	Immature teratoma
3	11	Oophorectomy	Right	Solid and cystic mass with haemorrhagic changes	22	Yolk sac tumor
4	24	Oophorectomy	Right	Solid mass	7	Yolk sac tumor
5	19	Oophorectomy	Left	Cyst with citrine content with intracystic vegetations	13	Mixed germ cell tumor
6	18	cystectomy	Right	Solid mass	3	Dysgerminoma
7	21	Oophorectomy	Right	Solid mass with haemorrhagic changes	20	Dysgerminoma
8	56	Hysterectomy adnexectomy	Bilateral	Solid mass and necrosis	18	Dysgerminoma
9	63	Oophorectomy	Right	Solid mass and hemorrhage	10	Dysgerminoma
10	46	Cyst biopsy	Unspecified	Poche kystique, sans végétation	2	Choriocarcinoma
11	54	Hysterectomy adnexectomy	Left	Tube-ovarian mass	1,5	Choriocarcinoma

Discussion

Germ cell tumors of the ovary are a heterogeneous group of tumors developed from primordial cells derived from the embryonic gonad. They are considered to be lesions linked to errors in the differentiation and/or migration of primordial cells [2]. The degree and the route of differentiation determine the histological type: the proliferation of undifferentiated germ cells induces a dysgerminoma, that of totipotent germ cells leads to an embryonic carcinoma, that of cells progressing in the extra-embryonic differentiation pathway leads to a tumor of the yolk sac or choriocarcinoma and that of cells with embryonic differentiation to mature and/or immature teratomas. Malignant germ cell tumors of the ovary (TGMO) are rare tumours. They represent 6% of ovarian tumours, with an annual incidence of around 0.5% per 100,000 women in France, i.e., around a hundred new cases annually [3]. During the study period, we recruited 68 cases of primary ovarian malignancy with 11 cases of germ cell tumor, i.e., 16.17% of cases. Over a 20-year period, Tewari et al found 72 cases of TGMO [4]. Mamouni N et al found 6 cases of TGMO in 4 years [5]. Malignant germ cell tumors of the ovary are linked to the existence of a cytogenetic characteristic common to all ovarian, testicular or extra-gonadal germ cell tumors, the presence of an isochromosome of the short arm of chromosome 12 [i(12p)] which is not found in

any other type of cancer [6]. These tumors occur at any age but are discovered most frequently between the 1st and 6th decades.

In the series by Mamouni N, et al. [5], the average age of the patients was 22 years with extremes of 1 and 30 years. For Tewari, et al. [4], the average age at diagnosis was 19 years, with extremes of 9 and 37 years [7]. In our series, the average age was higher with 30.45 years with extremes of 2 and 63 years, but in 64% of cases, our patients were under 25 years old. The revealing signs of an ovarian tumor are related to the consequences of the rapid increase in tumor volume leading to compression phenomena, pain which may be acute or chronic, an increase in abdominal volume with the perception of a mass, or more rarely torsion, hemorrhage or tumor rupture. A predominance of right-sided involvement is most often observed. In our series, it is above all the existence of an isolated abdominal mass or associated with abdominal pain or weight loss that has been reported. The right side was involved in 6 cases (54.54%), the left side in 2 cases. For 2 patients, the side was not specified, and one was bilateral. Macroscopically, the lesions differ according to histological type. Immature teratomas appear as a mass containing fat and solid portions with numerous microcystic structures of variable size [8]. Their diameter is between 9 and 28 cm with an average of 14 cm [9]. On the macroscopic level, for 2 of our cases, the masses were semi-solid, cerebroid in appearance

and semi-cystic with fatty content, measuring respectively 19 and 32 cm in long axis. They were larger compared to those reported in the literature, which can be explained by the rapid evolution of the tumor and the delay in consultation because sometimes people prefer to consult a traditional practitioner before seeing a doctor and the tumor has time to grow. In one of our cases, the patient underwent massage sessions which delayed the diagnosis and led to the extension of the tumor into the peritoneal cavity.

As for dysgerminoma, it presents as a solid, bulky tumor (on average 15 cm) with a bumpy outer surface and the section shows a lobulated, beige appearance containing necrotic and hemorrhagic cystic foci [10]. For our cases, the tumor measured between 3 and 20 cm long axis, with an average of 12.7 cm, which is comparable to what is reported in the literature. The same is true for the macroscopic aspect. For mixed germ cell tumours, the appearance depends on the type of lesion association present but most often they are solid in appearance [11]. In our case, the tumor was cystic with vegetations on its internal face. Histologically, there are 2 major groups of TGMO, dysgerminoma and non-germinomatous tumors. Dysgerminoma is the most frequent form with 30 to 40% of all TGMO and 10 to 15% are bilateral [12]. In our series, it is also the most frequent form (36.36%) with 1 case of bilaterality (9.09%). For Tewari, et al. [4] and Mamouni, et al. [5], it is in second place after immature teratomas. For non-germinomatous tumours, immature teratoma comes in 2nd position, with 20% of cases. It mainly affects young patients in the second decade [13]. In our series, we observed 2 cases of immature teratoma, 2 cases of Yolk sac tumor and 2 cases of non-gestational choriocarcinoma. In the series of Tewari, et al [4], the immature teratoma holds the 1st place (29 cases), followed by the mixed germ cell tumor (15 cases) and the yolk sac tumor (8 cases). For Mamouni N, et al. [5], immature teratoma also takes first place (3 cases), followed by mixed germ cell tumor (yolk sac tumor with dysgerminoma) and non-gestational choriocarcinoma.

Regarding therapeutic management, when the tumor affects a young woman of childbearing age, the surgeon should consider the possibility of a non-epithelial tumor and do a tumor marker assay before surgery for diagnostic orientation, in particular the assay of HCG and alpha fetoprotein. The gesture therefore consists at least of a unilateral annexectomy, a complete exploration of the pelvis and the entire abdominal cavity, peritoneal lavage and/or removal of any ascites present when the abdomen is opened, biopsies systematic peritoneal examinations (including at the level of the omentum) and a sample of any suspicious element. It is perfectly legitimate in young women in case of doubt, to perform the surgery in two stages to obtain the definitive results of the histology [14]. Rare malignant tumors of the ovary occur most often in young

women (median age: 20 years) even before the first pregnancy, and it is essential to be conservative and to respect the genital tract as much as possible to preserve fertility. In addition, they are tumors with a very good prognosis unlike epithelial tumors. For our cases, a radical treatment was carried out in 9 out of 11 cases because the tumors were discovered at an already advanced stage and required extensive excision.

Conclusion

TGMO are rare, we have only recruited 11 cases in 10 years. The most common histological type was dysgerminoma. TGMO are tumors that preferentially affect young women. Their growth is rapid with a significant increase in their volume, which makes conservative treatment difficult or impossible, thus hampering the obstetrical future of the patient. Earlier diagnosis would have improved management.

References

- Quirk JT, Natarajan N (2005) Ovarian cancer incidence in the United State, 1992-1999. *Gynecol Oncol* 97(2): 519-523.
- Surratt JT, Siegel MJ (1991) Imaging of pediatric ovarian masses. *Radiographics* 11: 533-548.
- Ray Coquard I, Pautier P, Pujade Lauraine E, Méeus P, Morice P, et al. (2010) Les tumeurs rares de l'ovaire : stratégies thérapeutiques en 2010, observatoire francophone des tumeurs rares de l'ovaires et émergence des centres de références. *Bull Cancer* 97(1) : 123-35.
- Tewari K, Cappuccini F, Disaia PJ, Berman ML, Manette A, et al. (2000) Malignant germ cell tumors of the ovary. *Obstetric and gynecology* 98(1): 128-133.
- Mamouni N, Saadi H, Erraghay S, Bouchikhi C, Banani A (2015) Tumeurs rares de l'ovaire : à propos de 11 cas de tumeurs non épithéliales malignes de l'ovaire. *Pan African Medical Journal* 20(174).
- Kurman RJ, Norris HJ (1976) Embryonal carcinoma of the ovary: a clinicopathologic entity distinct from endodermal sinus tumor resembling embryonal carcinoma of the adult testis. *Cancer* 38(6): 2420-2433.
- Gueye A, Narducci F, Baranzelli MC, Collinet P, Farine O, et al. (2007) Tumeurs germinales de l'ovaire. A propos de 36 cas. *Gynécologie Obstétrique & Fertilité* 35(5): 406-419.
- Yamaoka T, Togashi K, Koyama T, Fujiwara T, Higuchi T, et al. (2003) Immature teratoma of the ovary: correlation of MR imaging and pathologic findings *Eur Radiol* 13: 313-319.
- Kurman RJ, Norris HJ (1977) Malignant germ cell tumors of the ovary. *Hum Pathol* 8(5): 551-564.
- Tavassoli FA, Devilee P (2003) Tumors of the ovary and peritoneum. In: Tavassoli FA, Devilee P (Eds.), *World health organization of tumors pathology and genetics of tumors of the breast and female genital organs*. Lyon: IARC Press, pp.113-197.
- Brammer HM, Buck JL, Hayes WS, Sheth S, Tavassoli FA (1990) From the archives of the AFIP, malignant germ cell tumors of the ovary: radiologic-pathologic correlation. *Radiographics* 10(4): 715.
- Berek JS, Hacker NF (2005) Nonepithelial ovarian and fallopian tube cancers. In: Berek J, Hacker NF (Eds.), *Practical Gynecologic Oncology*. Philadelphia, PA: Lippincott Williams & Wilkins, pp. 511-541.

12. Gershenson DM, Del Junco G, Silva EG, Copeland, Larry J, et al. (1986) Immature teratoma of the ovary. *Obstet Gynecol* 68(5): 624-629.

13. Querleu D, Gladieff L, Delannes M, Mery E, Ferron G, et al. (2008) Preservation of fertility in gynaecologic cancers. *Bull Cancer* 95(5): 487-494.

ISSN: 2574-1241

DOI: 10.26717/BJSTR.2022.45.007212

HT Andrianjafitrimo. Biomed J Sci & Tech Res



This work is licensed under Creative Commons Attribution 4.0 License

Submission Link: <https://biomedres.us/submit-manuscript.php>



Assets of Publishing with us

- Global archiving of articles
- Immediate, unrestricted online access
- Rigorous Peer Review Process
- Authors Retain Copyrights
- Unique DOI for all articles

<https://biomedres.us/>